

2-12-1 血清総コレステロール値の分布(性・年齢階級別、服薬者を含む)

	総計		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上		
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
男性	総数	462	100.0	11	100.0	35	100.0	57	100.0	64	100.0	135	100.0	160	100.0	317	100.0	150	100.0	99	100.0
	120mg/d未満	5	1.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	5	3.1	0	0.0	0	0.0	5	5.1
	120-139	12	2.6	1	9.1	0	0.0	0	0.0	0	0.0	1	0.7	10	6.3	3	0.9	3	2.0	8	8.1
	140-159	42	9.1	3	27.3	0	0.0	4	7.0	2	3.1	11	8.1	22	13.8	24	7.6	15	10.0	15	15.2
	160-179	85	18.4	2	18.2	5	14.3	8	14.0	7	10.9	25	18.5	38	23.8	56	17.7	34	22.7	22	22.2
	180-199	109	23.6	1	9.1	12	34.3	13	22.8	19	29.7	35	25.9	29	18.1	82	25.9	40	26.7	14	14.1
	200-219	108	23.4	2	18.2	7	20.0	16	28.1	16	25.0	38	28.1	29	18.1	80	25.2	31	20.7	19	19.2
	220-239	57	12.3	1	9.1	6	17.1	11	19.3	15	23.4	10	7.4	14	8.8	41	12.9	11	7.3	9	9.1
	240-259	30	6.5	1	9.1	3	8.6	2	3.5	3	4.7	10	7.4	11	6.9	19	6.0	11	7.3	7	7.1
	260-279	9	1.9	0	0.0	2	5.7	3	5.3	1	1.6	2	1.5	1	0.6	7	2.2	2	1.3	0	0.0
	280-299	4	0.9	0	0.0	0	0.0	0	0.0	1	1.6	2	1.5	1	0.6	4	1.3	3	2.0	0	0.0
	300mg/d以上	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.7	0	0.0	1	0.3	0	0.0	0	0.0
	(再掲)220以上	101	21.9	2	18.2	11	31.4	16	28.1	20	31.3	25	18.5	27	16.9	72	22.7	27	18.0	16	16.2
	(再掲)240以上	44	9.5	1	9.1	5	14.3	5	8.8	5	7.8	15	11.1	13	8.1	31	9.8	16	10.7	7	7.1
	平均値(mg/d)	195.9		181.6		206.4		202.7		206.3		199.1		185.2		199.6		194.5		181.6	
	標準偏差	33.0		38.2		28.1		29.1		27.1		32.1		34.8		31.0		32.2		35.8	
女性	総数	543	100.0	19	100.0	45	100.0	78	100.0	84	100.0	153	100.0	164	100.0	376	100.0	148	100.0	103	100.0
	120mg/d未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	120-139	8	1.5	1	5.3	2	4.4	0	0.0	0	0.0	2	1.3	3	1.8	2	0.5	2	1.4	3	2.9
	140-159	32	5.9	3	15.8	5	11.1	8	10.3	1	1.2	4	2.6	11	6.7	16	4.3	6	4.1	8	7.8
	160-179	82	15.1	5	26.3	13	28.9	13	16.7	7	8.3	15	9.8	29	17.7	42	11.2	16	10.8	22	21.4
	180-199	125	23.0	4	21.1	12	26.7	20	25.6	16	19.0	28	18.3	45	27.4	82	21.8	35	23.6	27	26.2
	200-219	119	21.9	3	15.8	6	13.3	21	26.9	21	25.0	28	18.3	40	24.4	89	23.7	36	24.3	21	20.4
	220-239	79	14.5	1	5.3	6	13.3	4	5.1	18	21.4	33	21.6	17	10.4	63	16.8	24	16.2	9	8.7
	240-259	51	9.4	0	0.0	1	2.2	10	12.8	7	8.3	21	13.7	12	7.3	40	10.6	12	8.1	10	9.7
	260-279	28	5.2	1	5.3	0	0.0	1	1.3	10	11.9	11	7.2	5	3.0	25	6.6	8	5.4	2	1.9
	280-299	14	2.6	0	0.0	0	0.0	1	1.3	4	4.8	8	5.2	1	0.6	14	3.7	8	5.4	0	0.0
	300mg/d以上	5	0.9	1	5.3	0	0.0	0	0.0	0	0.0	3	2.0	1	0.6	3	0.8	1	0.7	1	1.0
	(再掲)220以上	177	32.6	3	15.8	7	15.6	16	20.5	39	46.4	76	49.7	36	22.0	145	38.6	53	35.8	22	21.4
	(再掲)240以上	98	18.0	2	10.5	1	2.2	12	15.4	21	25.0	43	28.1	19	11.6	82	21.8	29	19.6	13	12.6
	平均値(mg/d)	207.4		191.9		186.3		199.2		220.9		220.8		199.5		213.7		212.4		196.7	
	標準偏差	36.9		47.9		28.3		31.5		32.7		40.0		32.5		36.3		37.4		33.9	

注) コレステロールを下げる薬または中性脂肪(トリグリセライド)を下げる薬服用者を含む

2-12-2 血清総コレステロール値の分布(性・年齢階級別、服薬者除外)

	総計		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	389	100.0	10	100.0	34	100.0	55	100.0	56	100.0	109	100.0	125	100.0	266	100.0	115	100.0	79	100.0
120mg/dl未満	3	0.8	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	3	2.4	0	0.0	0	0.0	3	3.8
120-139	10	2.6	1	10.0	0	0.0	0	0.0	0	0.0	1	0.9	8	6.4	3	1.1	3	2.6	6	7.6
140-159	29	7.5	2	20.0	0	0.0	4	7.3	1	1.8	8	7.3	14	11.2	18	6.8	10	8.7	9	11.4
160-179	72	18.5	2	20.0	5	14.7	8	14.5	7	12.5	19	17.4	31	24.8	45	16.9	24	20.9	20	25.3
180-199	95	24.4	1	10.0	11	32.4	12	21.8	16	28.6	31	28.4	24	19.2	72	27.1	34	29.6	11	13.9
200-219	94	24.2	2	20.0	7	20.6	15	27.3	14	25.0	31	28.4	25	20.0	67	25.2	24	20.9	18	22.8
220-239	47	12.1	1	10.0	6	17.6	11	20.0	13	23.2	7	6.4	9	7.2	33	12.4	7	6.1	7	8.9
240-259	26	6.7	1	10.0	3	8.8	2	3.6	3	5.4	8	7.3	9	7.2	17	6.4	9	7.8	5	6.3
260-279	9	2.3	0	0.0	2	5.9	3	5.5	1	1.8	2	1.8	1	0.8	7	2.6	2	1.7	0	0.0
280-299	3	0.8	0	0.0	0	0.0	0	0.0	1	1.8	1	0.9	1	0.8	3	1.1	2	1.7	0	0.0
300mg/dl以上	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	1	0.9	0	0.0	1	0.4	0	0.0	0	0.0
(再掲)220以上	86	22.1	2	20.0	11	32.4	16	29.1	18	32.1	19	17.4	20	16.0	61	22.9	20	17.4	12	15.2
(再掲)240以上	39	10.0	1	10.0	5	14.7	5	9.1	5	8.9	12	11.0	11	8.8	28	10.5	13	11.3	5	6.3
平均値(mg/dl)	197.2		185.0		207.2		202.8		207.1		199.4		186.7		200.4		195.2		183.8	
標準偏差	32.4		38.5		28.2		29.5		27.0		31.6		34.2		30.9		31.8		34.1	
総数	420	100.0	19	100.0	45	100.0	75	100.0	71	100.0	113	100.0	97	100.0	297	100.0	100	100.0	59	100.0
120mg/dl未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
120-139	4	1.0	1	5.3	2	4.4	0	0.0	0	0.0	1	1.0	1	1.0	0	0.0	0	0.0	1	1.7
140-159	24	5.7	3	15.8	5	11.1	7	9.3	1	1.4	2	1.8	6	6.2	11	3.7	3	3.0	5	8.5
160-179	52	12.4	5	26.3	13	28.9	13	17.3	3	4.2	7	6.2	11	11.3	26	8.8	8	8.0	8	13.6
180-199	86	20.5	4	21.1	12	26.7	19	25.3	12	16.9	18	15.9	21	21.6	56	18.9	18	18.0	14	23.7
200-219	91	21.7	3	15.8	6	13.3	21	28.0	17	23.9	20	17.7	24	24.7	72	24.2	27	27.0	10	16.9
220-239	71	16.9	1	5.3	6	13.3	4	5.3	18	25.4	27	23.9	15	15.5	56	18.9	17	17.0	8	13.6
240-259	48	11.4	0	0.0	1	2.2	9	12.0	7	9.9	19	16.8	12	12.4	37	12.5	11	11.0	10	16.9
260-279	25	6.0	1	5.3	0	0.0	1	1.3	9	12.7	9	8.0	5	5.2	22	7.4	7	7.0	2	3.4
280-299	14	3.3	0	0.0	0	0.0	1	1.3	4	5.6	8	7.1	1	1.0	14	4.7	8	8.0	0	0.0
300mg/dl以上	5	1.2	1	5.3	0	0.0	0	0.0	0	0.0	3	2.7	1	1.0	3	1.0	1	1.0	1	1.7
(再掲)220以上	163	38.8	3	15.8	7	15.6	15	20.0	38	53.5	66	58.4	34	35.1	132	44.4	44	44.0	21	35.6
(再掲)240以上	92	21.9	2	10.5	1	2.2	11	14.7	20	28.2	39	34.5	19	19.6	76	25.6	27	27.0	13	22.0
平均値(mg/dl)	212.3		191.9		186.3		199.5		225.4		228.6		209.8		218.7		221.7		206.5	
標準偏差	37.9		47.9		28.3		31.1		31.9		39.6		34.4		36.4		37.2		37.2	

注) コレステロールを下げる薬または中性脂肪(トリグリセライド)を下げる薬服用者除外

2-13-1 血清HDL-コレステロール値の分布(性・年齢階級別、服薬者を含む)

	総計		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上		
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
男性	総数	462	100.0	11	100.0	35	100.0	57	100.0	64	100.0	135	100.0	160	100.0	317	100.0	150	100.0	99	100.0
	30mg/dl未満	4	0.9	0	0.0	0	0.0	0	0.0	0	0.0	1	0.7	3	1.9	2	0.6	1	0.7	2	2.0
	30-39	52	11.3	0	0.0	1	2.9	5	8.8	6	9.4	15	11.1	25	15.6	40	12.6	24	16.0	11	11.1
	40-49	112	24.2	3	27.3	10	28.6	16	28.1	12	18.8	30	22.2	41	25.6	73	23.0	32	21.3	26	26.3
	50-59	125	27.1	4	36.4	11	31.4	13	22.8	19	29.7	31	23.0	47	29.4	79	24.9	37	24.7	31	31.3
	60-69	92	19.9	3	27.3	8	22.9	14	24.6	11	17.2	34	25.2	22	13.8	65	20.5	29	19.3	16	16.2
	70-79	45	9.7	0	0.0	3	8.6	3	5.3	9	14.1	14	10.4	16	10.0	31	9.8	17	11.3	11	11.1
	80-89	22	4.8	1	9.1	0	0.0	4	7.0	4	6.3	8	5.9	5	3.1	20	6.3	10	6.7	1	1.0
	90-99	7	1.5	0	0.0	2	5.7	1	1.8	2	3.1	1	0.7	1	0.6	4	1.3	0	0.0	1	1.0
	100mg/dl以上	3	0.6	0	0.0	0	0.0	1	1.8	1	1.6	1	0.7	0	0.0	3	0.9	0	0.0	0	0.0
	平均値(mg/dl)	56.2		56.6		57.6		56.9		59.9		57.4		53.1		56.7		55.3		54.1	
	標準偏差	14.9		11.3		13.6		14.7		16.7		15.5		13.8		15.6		14.8		13.2	
女性	総数	543	100.0	19	100.0	45	100.0	78	100.0	84	100.0	153	100.0	164	100.0	376	100.0	148	100.0	103	100.0
	30mg/dl未満	2	0.4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	1.2	0	0.0	0	0.0	2	1.9
	30-39	18	3.3	0	0.0	1	2.2	1	1.3	2	2.4	7	4.6	7	4.3	13	3.5	6	4.1	4	3.9
	40-49	66	12.2	1	5.3	3	6.7	6	7.7	8	9.5	15	9.8	33	20.1	38	10.1	18	12.2	24	23.3
	50-59	123	22.7	5	26.3	8	17.8	12	15.4	14	16.7	41	26.8	43	26.2	85	22.6	45	30.4	25	24.3
	60-69	137	25.2	9	47.4	15	33.3	21	26.9	19	22.6	31	20.3	42	25.6	89	23.7	36	24.3	24	23.3
	70-79	107	19.7	2	10.5	11	24.4	21	26.9	17	20.2	32	20.9	24	14.6	78	20.7	23	15.5	16	15.5
	80-89	50	9.2	2	10.5	4	8.9	11	14.1	12	14.3	13	8.5	8	4.9	39	10.4	10	6.8	5	4.9
	90-99	26	4.8	0	0.0	2	4.4	3	3.8	7	8.3	9	5.9	5	3.0	21	5.6	7	4.7	3	2.9
	100mg/dl以上	14	2.6	0	0.0	1	2.2	3	3.8	5	6.0	5	3.3	0	0.0	13	3.5	3	2.0	0	0.0
	平均値(mg/dl)	64.9		64.6		67.0		68.9		70.1		65.5		59.3		66.4		63.2		58.7	
	標準偏差	16.0		10.5		15.5		15.0		17.6		16.6		14.0		16.2		15.6		14.6	

注)コレステロールを下げる薬または中性脂肪(トリグリセライド)を下げる薬服用者を含む

2-13-2 血清HDL-コレステロール値の分布(性・年齢階級別、服薬者除外)

	総計		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上		
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
男性	総数	389	100.0	10	100.0	34	100.0	55	100.0	56	100.0	109	100.0	125	100.0	266	100.0	115	100.0	79	100.0
	30mg/dl未満	4	1.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.9	3	2.4	2	0.8	1	0.9	2	2.5
	30-39	47	12.1	0	0.0	0	0.0	5	9.1	6	10.7	13	11.9	23	18.4	37	13.9	21	18.3	10	12.7
	40-49	98	25.2	3	30.0	10	29.4	15	27.3	12	21.4	25	22.9	33	26.4	63	23.7	24	20.9	22	27.8
	50-59	90	23.1	3	30.0	11	32.4	12	21.8	12	21.4	23	21.1	29	23.2	57	21.4	24	20.9	19	24.1
	60-69	81	20.8	3	30.0	8	23.5	14	25.5	10	17.9	27	24.8	19	15.2	56	21.1	24	20.9	14	17.7
	70-79	40	10.3	0	0.0	3	8.8	3	5.5	9	16.1	12	11.0	13	10.4	27	10.2	13	11.3	10	12.7
	80-89	20	5.1	1	10.0	0	0.0	4	7.3	4	7.1	7	6.4	4	3.2	18	6.8	8	7.0	1	1.3
	90-99	6	1.5	0	0.0	2	5.9	1	1.8	2	3.6	0	0.0	1	0.8	3	1.1	0	0.0	1	1.3
	100mg/dl以上	3	0.8	0	0.0	0	0.0	1	1.8	1	1.8	1	0.9	0	0.0	3	1.1	0	0.0	0	0.0
	平均値(mg/dl)	56.3		57.2		58.2		57.3		60.5		57.3		52.6		56.7		55.0		54.2	
	標準偏差	15.4		11.7		13.2		14.9		17.7		15.8		14.6		16.1		15.3		14.3	
女性	総数	420	100.0	19	100.0	45	100.0	75	100.0	71	100.0	113	100.0	97	100.0	297	100.0	100	100.0	59	100.0
	30mg/dl未満	2	0.5	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	2.1	0	0.0	0	0.0	2	3.4
	30-39	13	3.1	0	0.0	1	2.2	1	1.3	2	2.8	5	4.4	4	4.1	10	3.4	3	3.0	2	3.4
	40-49	47	11.2	1	5.3	3	6.7	5	6.7	7	9.9	11	9.7	20	20.6	30	10.1	14	14.0	13	22.0
	50-59	89	21.2	5	26.3	8	17.8	11	14.7	11	15.5	30	26.5	24	24.7	60	20.2	29	29.0	16	27.1
	60-69	105	25.0	9	47.4	15	33.3	21	28.0	16	22.5	22	19.5	22	22.7	71	23.9	24	24.0	10	16.9
	70-79	86	20.5	2	10.5	11	24.4	20	26.7	14	19.7	25	22.1	14	14.4	63	21.2	15	15.0	10	16.9
	80-89	43	10.2	2	10.5	4	8.9	11	14.7	10	14.1	9	8.0	7	7.2	33	11.1	8	8.0	4	6.8
	90-99	21	5.0	0	0.0	2	4.4	3	4.0	6	8.5	6	5.3	4	4.1	17	5.7	4	4.0	2	3.4
	100mg/dl以上	14	3.3	0	0.0	1	2.2	3	4.0	5	7.0	5	4.4	0	0.0	13	4.4	3	3.0	0	0.0
	平均値(mg/dl)	66.0		64.6		67.0		69.5		70.7		65.9		59.9		67.4		63.8		58.8	
	標準偏差	16.3		10.5		15.5		14.8		18.1		16.9		15.1		16.6		16.0		15.7	

注)コレステロールを下げる薬または中性脂肪(トリグリセライド)を下げる薬服用者除外

2-14-1 血清LDL-コレステロール値の分布(性・年齢階級別、服薬者を含む)

	総計		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	482	100.0	11	100.0	35	100.0	57	100.0	64	100.0	135	100.0	160	100.0	317	100.0	150	100.0	99	100.0
40mg/dl未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
40-49	3	0.6	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	3	1.9	0	0.0	0	0.0	3	3.0
50-59	7	1.5	0	0.0	0	0.0	1	1.8	1	1.6	3	2.2	2	1.3	5	1.6	2	1.3	2	2.0
60-69	20	4.3	0	0.0	0	0.0	1	1.8	2	3.1	6	4.4	11	6.9	11	3.5	7	4.7	9	9.1
70-79	34	7.4	2	18.2	1	2.9	3	5.3	1	1.6	8	5.9	19	11.9	21	6.6	17	11.3	10	10.1
80-89	47	10.2	1	9.1	3	8.6	6	10.5	3	4.7	16	11.9	18	11.3	31	9.8	16	10.7	12	12.1
90-99	47	10.2	2	18.2	1	2.9	2	3.5	6	9.4	14	10.4	22	13.8	34	10.7	21	14.0	10	10.1
100-109	63	13.6	2	18.2	6	17.1	4	7.0	10	15.6	23	17.0	18	11.3	41	12.9	21	14.0	14	14.1
110-119	52	11.3	1	9.1	5	14.3	3	5.3	11	17.2	19	14.1	13	8.1	38	12.0	17	11.3	8	8.1
120-129	68	14.7	1	9.1	5	14.3	11	19.3	11	17.2	19	14.1	21	13.1	49	15.5	18	12.0	13	13.1
130-139	42	9.1	0	0.0	6	17.1	9	15.8	5	7.8	7	5.2	15	9.4	25	7.9	7	4.7	11	11.1
140-149	31	6.7	0	0.0	1	2.9	7	12.3	5	7.8	10	7.4	8	5.0	29	9.1	13	8.7	1	1.0
150-159	21	4.5	1	9.1	3	8.6	7	12.3	4	6.3	2	1.5	4	2.5	15	4.7	4	2.7	2	2.0
160-169	16	3.5	0	0.0	3	8.6	2	3.5	2	3.1	4	3.0	5	3.1	9	2.8	3	2.0	4	4.0
170-179	6	1.3	1	9.1	1	2.9	0	0.0	3	4.7	1	0.7	0	0.0	4	1.3	1	0.7	0	0.0
180-189	3	0.6	0	0.0	0	0.0	1	1.8	0	0.0	2	1.5	0	0.0	3	0.9	2	1.3	0	0.0
190-199	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
200-209	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
210-219	2	0.4	0	0.0	0	0.0	0	0.0	0	0.0	1	0.7	1	0.6	2	0.6	1	0.7	0	0.0
220mg/dl以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
平均値(mg/dl)	112.1		108.4		124.2		122.4		119.0		110.5		104.7		114.0		108.2		102.3	
標準偏差	28.9		32.7		25.6		28.8		26.6		28.1		28.8		28.5		28.4		28.5	
総数	543	100.0	19	100.0	45	100.0	78	100.0	84	100.0	153	100.0	164	100.0	376	100.0	148	100.0	103	100.0
40mg/dl未満	1	0.2	0	0.0	0	0.0	0	0.0	1	1.2	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0
40-49	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.6	0	0.0	0	0.0	1	1.0
50-59	7	1.3	1	5.3	2	4.4	1	1.3	0	0.0	1	0.7	2	1.2	2	0.5	1	0.7	2	1.9
60-69	12	2.2	0	0.0	1	2.2	4	5.1	1	1.2	1	0.7	5	3.0	8	2.1	3	2.0	3	2.9
70-79	33	6.1	1	5.3	5	11.1	6	7.7	2	2.4	5	3.3	14	8.5	16	4.3	7	4.7	11	10.7
80-89	47	8.7	3	15.8	4	8.9	5	6.4	6	7.1	11	7.2	18	11.0	28	7.4	12	8.1	12	11.7
90-99	62	11.4	5	26.3	11	24.4	8	10.3	3	3.6	9	5.9	26	15.9	30	8.0	15	10.1	16	15.5
100-109	76	14.0	2	10.5	5	11.1	14	17.9	12	14.3	20	13.1	23	14.0	53	14.1	20	13.5	16	15.5
110-119	67	12.3	1	5.3	7	15.6	11	14.1	11	13.1	17	11.1	20	12.2	47	12.5	16	10.8	12	11.7
120-129	67	12.3	2	10.5	3	6.7	14	17.9	9	10.7	20	13.1	19	11.6	53	14.1	20	13.5	9	8.7
130-139	51	9.4	1	5.3	3	6.7	2	2.6	13	15.5	20	13.1	12	7.3	41	10.9	18	12.2	6	5.8
140-149	45	8.3	1	5.3	1	2.2	6	7.7	10	11.9	19	12.4	8	4.9	39	10.4	13	8.8	4	3.9
150-159	29	5.3	0	0.0	1	2.2	3	3.8	5	6.0	10	6.5	10	6.1	21	5.6	9	6.1	7	6.8
160-169	18	3.3	0	0.0	2	4.4	3	3.8	3	3.6	7	4.6	3	1.8	13	3.5	3	2.0	3	2.9
170-179	8	1.5	0	0.0	0	0.0	1	1.3	3	3.6	3	2.0	1	0.6	8	2.1	4	2.7	0	0.0
180-189	5	0.9	1	5.3	0	0.0	0	0.0	3	3.6	1	0.7	0	0.0	4	1.1	1	0.7	0	0.0
190-199	5	0.9	0	0.0	0	0.0	0	0.0	1	1.2	4	2.6	0	0.0	5	1.3	2	1.4	0	0.0
200-209	5	0.9	0	0.0	0	0.0	0	0.0	1	1.2	2	1.3	2	1.2	4	1.1	2	1.4	1	1.0
210-219	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.7	0	0.0	1	0.3	1	0.7	0	0.0
220mg/dl以上	3	0.6	1	5.3	0	0.0	0	0.0	0	0.0	2	1.3	0	0.0	2	0.5	1	0.7	0	0.0
平均値(mg/dl)	116.7		110.6		103.0		110.3		126.2		127.3		109.4		121.3		120.7		107.0	
標準偏差	31.7		40.8		25.5		26.7		30.9		34.1		28.0		31.6		32.3		28.4	

注)コレステロールを下げる薬または中性脂肪(トリグリセリド)を下げる薬服用者を含む

2-14-2 血清LDL-コレステロール値の分布(性・年齢階級別、服薬者除外)

	総計		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	389	100.0	10	100.0	34	100.0	55	100.0	56	100.0	109	100.0	125	100.0	266	100.0	115	100.0	79	100.0
40mg/dl未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
40-49	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.8	0	0.0	0	0.0	0	0.0
50-59	5	1.3	0	0.0	0	0.0	1	1.8	1	1.8	2	1.8	1	0.8	4	1.5	1	0.9	1	1.3
60-69	16	4.1	0	0.0	0	0.0	1	1.8	1	1.8	4	3.7	10	8.0	8	3.0	5	4.3	8	10.1
70-79	25	6.4	2	20.0	1	2.9	3	5.5	0	0.0	8	7.3	11	8.8	16	6.0	13	11.3	6	7.6
80-89	37	9.5	1	10.0	3	8.8	6	10.9	2	3.6	13	11.9	12	9.6	24	9.0	10	8.7	9	11.4
90-99	38	9.8	1	10.0	1	2.9	2	3.6	6	10.7	10	9.2	18	14.4	28	10.5	16	13.9	8	10.1
100-109	53	13.6	2	20.0	6	17.6	4	7.3	8	14.3	16	14.7	17	13.6	31	11.7	15	13.0	14	17.7
110-119	46	11.8	1	10.0	4	11.8	3	5.5	11	19.6	17	15.6	10	8.0	35	13.2	15	13.0	6	7.6
120-129	84	16.5	1	10.0	5	14.7	10	18.2	11	19.6	19	17.4	18	14.4	46	17.3	16	13.9	12	15.2
130-139	35	9.0	0	0.0	6	17.6	9	16.4	4	7.1	5	4.6	11	8.8	21	7.9	5	4.3	8	10.1
140-149	26	6.7	0	0.0	1	2.9	7	12.7	3	5.4	8	7.3	7	5.6	24	9.0	11	9.6	1	1.3
150-159	18	4.6	1	10.0	3	8.8	6	10.9	4	7.1	1	0.9	3	2.4	13	4.9	3	2.6	1	1.3
160-169	15	3.9	0	0.0	3	8.8	2	3.6	2	3.6	3	2.8	5	4.0	8	3.0	2	1.7	4	5.1
170-179	6	1.5	1	10.0	1	2.9	0	0.0	3	5.4	1	0.9	0	0.0	4	1.5	1	0.9	0	0.0
180-189	2	0.5	0	0.0	0	0.0	1	1.8	0	0.0	1	0.9	0	0.0	2	0.8	1	0.9	0	0.0
190-199	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
200-209	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
210-219	2	0.5	0	0.0	0	0.0	0	0.0	0	0.0	1	0.9	1	0.8	2	0.8	1	0.9	0	0.0
220mg/dl以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
平均値(mg/dl)	113.9		110.2		124.4		122.0		120.7		111.0		107.2		115.5		109.7		104.5	
標準偏差	28.4		33.9		26.0		29.1		25.6		27.4		28.4		28.2		27.9		27.1	
総数	420	100.0	19	100.0	45	100.0	75	100.0	71	100.0	113	100.0	97	100.0	297	100.0	100	100.0	59	100.0
40mg/dl未満	1	0.2	0	0.0	0	0.0	0	0.0	1	1.4	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0
40-49	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	1.0	0	0.0	0	0.0	1	1.7
50-59	6	1.4	1	5.3	2	4.4	1	1.3	0	0.0	1	0.9	1	1.0	2	0.7	1	1.0	1	1.7
60-69	7	1.7	0	0.0	1	2.2	4	5.3	1	1.4	1	0.9	0	0.0	6	2.0	1	1.0	0	0.0
70-79	21	5.0	1	5.3	5	11.1	5	6.7	2	2.8	1	0.9	7	7.2	10	3.4	3	3.0	5	8.5
80-89	24	5.7	3	15.8	4	8.9	5	6.7	1	1.4	4	3.5	7	7.2	11	3.7	2	2.0	6	10.2
90-99	38	9.0	5	26.3	11	24.4	8	10.7	2	2.8	5	4.4	7	7.2	17	5.7	5	5.0	5	8.5
100-109	61	14.5	2	10.5	5	11.1	14	18.7	9	12.7	16	14.2	15	15.5	45	15.2	17	17.0	9	15.3
110-119	47	11.2	1	5.3	7	15.6	10	13.3	9	12.7	9	8.0	11	11.3	33	11.1	8	8.0	6	10.2
120-129	60	14.3	2	10.5	3	6.7	14	18.7	9	12.7	16	14.2	16	16.5	48	16.2	17	17.0	7	11.9
130-139	44	10.5	1	5.3	3	6.7	2	2.7	12	16.9	17	15.0	9	9.3	35	11.8	14	14.0	5	8.5
140-149	42	10.0	1	5.3	1	2.2	6	8.0	10	14.1	16	14.2	8	8.2	36	12.1	10	10.0	4	6.8
150-159	26	6.2	0	0.0	1	2.2	2	2.7	5	7.0	9	8.0	9	9.3	19	6.4	9	9.0	6	10.2
160-169	16	3.8	0	0.0	2	4.4	3	4.0	3	4.2	5	4.4	3	3.1	11	3.7	2	2.0	3	5.1
170-179	8	1.9	0	0.0	0	0.0	1	1.3	3	4.2	3	2.7	1	1.0	8	2.7	4	4.0	0	0.0
180-189	4	1.0	1	5.3	0	0.0	0	0.0	2	2.8	1	0.9	0	0.0	3	1.0	1	1.0	0	0.0
190-199	5	1.2	0	0.0	0	0.0	0	0.0	1	1.4	4	3.5	0	0.0	5	1.7	2	2.0	0	0.0
200-209	5	1.2	0	0.0	0	0.0	0	0.0	1	1.4	2	1.8	2	2.1	4	1.3	2	2.0	1	1.7
210-219	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.9	0	0.0	1	0.3	1	1.0	0	0.0
220mg/dl以上	3	0.7	1	5.3	0	0.0	0	0.0	0	0.0	2	1.8	0	0.0	2	0.7	1	1.0	0	0.0
平均値(mg/dl)	121.1		110.6		103.0		110.2		129.6		133.5		119.1		125.5		129.4		115.9	
標準偏差	32.4		40.6		25.5		26.4		30.3		34.9		29.1		32.0		32.7		30.5	

注)コレステロールを下げる薬または中性脂肪(トリグリセリド)を下げる薬服用者除外

2-15 メタボリックシンドローム(内臓脂肪症候群)の状況(性・年齢階級別)

	総計		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上		
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
総数	995	100.0	26	100.0	78	100.0	134	100.0	147	100.0	288	100.0	322	100.0	690	100.0	297	100.0	201	100.0	
総数	169	17.0	1	3.8	1	1.3	8	6.0	16	10.9	59	20.5	84	26.1	118	17.1	77	25.9	49	24.4	
メタボリックシンドローム(内臓脂肪症候群)が強く疑われる者	171	17.2	2	7.7	7	9.0	17	12.7	34	23.1	51	17.7	60	18.6	123	17.8	50	16.8	39	19.4	
メタボリックシンドローム(内臓脂肪症候群)の予備群と考えられる者	655	65.8	23	88.5	70	89.7	109	81.3	97	66.0	178	61.8	178	55.3	449	65.1	170	57.2	113	56.2	
上記以外																					
総数	461	100.0	10	100.0	35	100.0	57	100.0	64	100.0	135	100.0	160	100.0	317	100.0	150	100.0	99	100.0	
総数	120	26.0	1	10.0	1	2.9	5	8.8	11	17.2	45	33.3	57	35.6	88	27.8	60	40.0	30	30.3	
メタボリックシンドローム(内臓脂肪症候群)が強く疑われる者	121	26.2	2	20.0	5	14.3	14	24.6	23	35.9	39	28.9	38	23.8	86	27.1	33	22.0	28	28.3	
メタボリックシンドローム(内臓脂肪症候群)の予備群と考えられる者	220	47.7	7	70.0	29	82.9	38	66.7	30	46.9	51	37.8	65	40.6	143	45.1	57	38.0	41	41.4	
上記以外																					
総数	534	100.0	16	100.0	43	100.0	77	100.0	83	100.0	153	100.0	162	100.0	373	100.0	147	100.0	102	100.0	
総数	49	9.2	0	0.0	0	0.0	3	3.9	5	6.0	14	9.2	27	16.7	30	8.0	17	11.6	19	18.6	
メタボリックシンドローム(内臓脂肪症候群)が強く疑われる者	50	9.4	0	0.0	2	4.7	3	3.9	11	13.3	12	7.8	22	13.6	37	9.9	17	11.6	11	10.8	
メタボリックシンドローム(内臓脂肪症候群)の予備群と考えられる者	435	81.5	16	100.0	41	95.3	71	92.2	67	80.7	127	83.0	113	69.8	306	82.0	113	76.9	72	70.6	
上記以外																					

注1) 血圧、糖質、ヘモグロビンA1c、血清HDLコレステロール値の測定を行い、身体状況調査の問題において血圧を下げる薬、インスリン注射または血糖を下げる薬、コレステロールを下げる薬、中性脂肪(トリグリセライド)を下げる薬の服用状況に全て回答した20歳以上の者を集計対象とした。なお、女性に妊婦を除外した。
注2) 判定基準はP13参照。

2-16 腹囲区別、血中脂質、血圧、血糖のいずれかのリスクを有する割合(性・年齢階級別)

	腹囲	総数		リスク保有数								
				0		1つ		2つ		3つ		
		人数	%	人数	%	人数	%	人数	%	人数	%	
総数	総数	<85cm(男)/90cm(女)	603	100.0	242	40.1	231	38.3	97	16.1	33	5.5
		≥85cm(男)/90cm(女)	392	100.0	52	13.3	171	43.6	118	30.1	51	13.0
	20-29歳	<85cm(男)/90cm(女)	20	100.0	19	95.0	1	5.0	0	0.0	0	0.0
		≥85cm(男)/90cm(女)	6	100.0	3	50.0	2	33.3	1	16.7	0	0.0
	30-39歳	<85cm(男)/90cm(女)	58	100.0	46	79.3	12	20.7	0	0.0	0	0.0
		≥85cm(男)/90cm(女)	20	100.0	12	60.0	7	35.0	0	0.0	1	5.0
	40-49歳	<85cm(男)/90cm(女)	95	100.0	74	77.9	20	21.1	1	1.1	0	0.0
		≥85cm(男)/90cm(女)	39	100.0	14	35.9	17	43.6	6	15.4	2	5.1
	50-59歳	<85cm(男)/90cm(女)	90	100.0	37	41.1	42	46.7	9	10.0	2	2.2
		≥85cm(男)/90cm(女)	57	100.0	7	12.3	34	59.6	10	17.5	6	10.5
	60-69歳	<85cm(男)/90cm(女)	166	100.0	44	26.5	77	46.4	35	21.1	10	6.0
		≥85cm(男)/90cm(女)	122	100.0	12	9.8	51	41.8	40	32.8	19	15.6
	70歳以上	<85cm(男)/90cm(女)	174	100.0	22	12.6	79	45.4	52	29.9	21	12.1
		≥85cm(男)/90cm(女)	148	100.0	4	2.7	60	40.5	61	41.2	23	15.5
(再掲)40-74歳	<85cm(男)/90cm(女)	414	100.0	166	40.1	170	41.1	60	14.5	18	4.3	
	≥85cm(男)/90cm(女)	276	100.0	35	12.7	123	44.6	81	29.3	37	13.4	
(再掲)65-74歳	<85cm(男)/90cm(女)	164	100.0	31	18.9	85	51.8	37	22.6	11	6.7	
	≥85cm(男)/90cm(女)	133	100.0	6	4.5	50	37.6	55	41.4	22	16.5	
(再掲)75歳以上	<85cm(男)/90cm(女)	111	100.0	11	9.9	48	43.2	37	33.3	15	13.5	
	≥85cm(男)/90cm(女)	90	100.0	2	2.2	39	43.3	36	40.0	13	14.4	
男性	総数	<85cm	182	100.0	64	35.2	85	46.7	29	15.9	4	2.2
		≥85cm	279	100.0	38	13.6	121	43.4	87	31.2	33	11.8
	20-29歳	<85cm	4	100.0	3	75.0	1	25.0	0	0.0	0	0.0
		≥85cm	6	100.0	3	50.0	2	33.3	1	16.7	0	0.0
	30-39歳	<85cm	21	100.0	14	66.7	7	33.3	0	0.0	0	0.0
		≥85cm	14	100.0	8	57.1	5	35.7	0	0.0	1	7.1
	40-49歳	<85cm	29	100.0	20	69.0	9	31.0	0	0.0	0	0.0
		≥85cm	28	100.0	9	32.1	14	50.0	5	17.9	0	0.0
	50-59歳	<85cm	26	100.0	8	30.8	14	53.8	4	15.4	0	0.0
		≥85cm	38	100.0	4	10.5	23	60.5	7	18.4	4	10.5
	60-69歳	<85cm	40	100.0	10	25.0	21	52.5	8	20.0	1	2.5
		≥85cm	95	100.0	11	11.6	39	41.1	33	34.7	12	12.6
	70歳以上	<85cm	62	100.0	9	14.5	33	53.2	17	27.4	3	4.8
		≥85cm	98	100.0	3	3.1	38	38.8	41	41.8	16	16.3
(再掲)40-74歳	<85cm	118	100.0	42	35.6	57	48.3	18	15.3	1	0.8	
	≥85cm	199	100.0	25	12.6	86	43.2	63	31.7	25	12.6	
(再掲)65-74歳	<85cm	52	100.0	12	23.1	29	55.8	10	19.2	1	1.9	
	≥85cm	98	100.0	5	5.1	33	33.7	43	43.9	17	17.3	
(再掲)75歳以上	<85cm	39	100.0	5	12.8	20	51.3	11	28.2	3	7.7	
	≥85cm	60	100.0	2	3.3	28	46.7	23	38.3	7	11.7	
女性	総数	<90cm	421	100.0	178	42.3	146	34.7	68	16.2	29	6.9
		≥90cm	113	100.0	14	12.4	50	44.2	31	27.4	18	15.9
	20-29歳	<90cm	16	100.0	16	100.0	0	0.0	0	0.0	0	0.0
		≥90cm	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	30-39歳	<90cm	37	100.0	32	86.5	5	13.5	0	0.0	0	0.0
		≥90cm	6	100.0	4	66.7	2	33.3	0	0.0	0	0.0
	40-49歳	<90cm	66	100.0	54	81.8	11	16.7	1	1.5	0	0.0
		≥90cm	11	100.0	5	45.5	3	27.3	1	9.1	2	18.2
	50-59歳	<90cm	64	100.0	29	45.3	28	43.8	5	7.8	2	3.1
		≥90cm	19	100.0	3	15.8	11	57.9	3	15.8	2	10.5
	60-69歳	<90cm	126	100.0	34	27.0	56	44.4	27	21.4	9	7.1
		≥90cm	27	100.0	1	3.7	12	44.4	7	25.9	7	25.9
	70歳以上	<90cm	112	100.0	13	11.6	46	41.1	35	31.3	18	16.1
		≥90cm	50	100.0	1	2.0	22	44.0	20	40.0	7	14.0
(再掲)40-74歳	<90cm	296	100.0	124	41.9	113	38.2	42	14.2	17	5.7	
	≥90cm	77	100.0	10	13.0	37	48.1	18	23.4	12	15.6	
(再掲)65-74歳	<90cm	112	100.0	19	17.0	56	50.0	27	24.1	10	8.9	
	≥90cm	35	100.0	1	2.9	17	48.6	12	34.3	5	14.3	
(再掲)75歳以上	<90cm	72	100.0	6	8.3	28	38.9	26	36.1	12	16.7	
	≥90cm	30	100.0	0	0.0	11	36.7	13	43.3	6	20.0	

注1) 血圧、腹囲、ヘモグロビンA1c、血清HDLコレステロール値の測定を行い、身体状況調査の間診において血圧を下げる薬、インスリン注射または血糖を下げる薬、コレステロールを下げる薬、中性脂肪(トリグリセリド)を下げる薬の服用状況に全て回答した20歳以上の者を集計対象とした。なお、女性は妊婦を除外した。
 注2) リスク項目の基準はp.13参照

2-17 「糖尿病が強く疑われる人」及び「糖尿病の可能性が否定できない人」の状況(性・年齢級別)

	総計		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	901	100.0	27	100.0	68	100.0	121	100.0	131	100.0	266	100.0	288	100.0	631	100.0	277	100.0	175	100.0
糖尿病が強く疑われる人	106	11.8	0	0.0	1	1.5	3	2.5	9	6.9	35	13.2	58	20.1	66	10.5	42	15.2	39	22.3
<small>(再掲)糖尿病が強く疑われる人のうち服薬者※</small>	69	65.1	0	0.0	0	0.0	2	66.7	6	66.7	22	62.9	39	67.2	41	62.1	25	59.5	28	71.8
糖尿病の可能性が否定できない人	80	8.9	0	0.0	2	2.9	5	4.1	8	6.1	32	12.0	33	11.5	55	8.7	33	11.9	23	13.1
上記以外	715	79.4	27	100.0	65	95.6	113	93.4	114	87.0	199	74.8	197	68.4	510	80.8	202	72.9	113	64.6
総数	418	100.0	10	100.0	31	100.0	50	100.0	57	100.0	123	100.0	147	100.0	286	100.0	138	100.0	91	100.0
糖尿病が強く疑われる人	61	14.6	0	0.0	1	3.2	2	4.0	3	5.3	20	16.3	35	23.8	37	12.9	24	17.4	23	25.3
<small>(再掲)糖尿病が強く疑われる人のうち服薬者※</small>	40	65.6	0	0.0	0	0.0	1	50.0	2	66.7	14	70.0	23	65.7	24	64.9	15	62.5	16	69.6
糖尿病の可能性が否定できない人	37	8.9	0	0.0	0	0.0	1	2.0	6	10.5	14	11.4	16	10.9	27	9.4	18	13.0	10	11.0
上記以外	320	76.6	10	100.0	30	96.8	47	94.0	48	84.2	89	72.4	96	65.3	222	77.6	96	69.6	58	63.7
総数	483	100.0	17	100.0	37	100.0	71	100.0	74	100.0	143	100.0	141	100.0	345	100.0	139	100.0	84	100.0
糖尿病が強く疑われる人	45	9.3	0	0.0	0	0.0	1	1.4	6	8.1	15	10.5	23	16.3	29	8.4	18	12.9	16	19.0
<small>(再掲)糖尿病が強く疑われる人のうち服薬者※</small>	29	64.4	0	0.0	0	0.0	1	100.0	4	66.7	8	53.3	16	69.6	17	58.6	10	55.6	12	75.0
糖尿病の可能性が否定できない人	43	8.9	0	0.0	2	5.4	4	5.6	2	2.7	18	12.6	17	12.1	28	8.1	15	10.8	13	15.5
上記以外	395	81.8	17	100.0	35	94.6	66	93.0	66	89.2	110	76.9	101	71.6	288	83.5	106	76.3	55	65.5

注1) 糖尿病が強く疑われる人は、ヘモグロビンA1cの値が6.5%以上の者、または「インスリン注射または血糖を下げる薬」を服用している」と回答した者、または健康意識調査で「糖尿病の治療を受けている」と回答した者とした。
 糖尿病の可能性が否定できない人は、ヘモグロビンA1cの値が6.0%以上6.5%未満の者で、糖尿病が強く疑われる者以外の者とした。

注2) 「服薬者」とは、質問票で「インスリン注射または血糖値を下げる薬」の使用有と回答した者

2-18 高血圧症有病者の状況(性・年齢階級別)

	総計		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	1,146	100.0	38	100.0	86	100.0	148	100.0	163	100.0	332	100.0	379	100.0	787	100.0	348	100.0	235	100.0
高血圧症有病者	569	49.7	2	5.3	7	8.1	23	15.5	69	42.3	202	60.8	266	70.2	390	49.6	227	65.2	170	72.3
(再掲)高血圧症有病者のうち服薬者※	375	65.9	0	0.0	1	14.3	6	26.1	34	49.3	125	61.9	209	78.6	237	60.8	153	67.4	137	80.6
正常高値血圧者	255	22.3	5	13.2	10	11.6	19	12.8	42	25.8	74	22.3	105	27.7	178	22.6	96	27.6	62	26.4
上記以外	322	28.1	31	81.6	69	80.2	106	71.6	52	31.9	56	16.9	8	2.1	219	27.8	25	7.2	3	1.3
総数	520	100.0	14	100.0	36	100.0	62	100.0	71	100.0	152	100.0	185	100.0	355	100.0	169	100.0	115	100.0
高血圧症有病者	297	57.1	2	14.3	6	16.7	17	27.4	37	52.1	102	67.1	133	71.9	208	58.6	119	70.4	81	70.4
(再掲)高血圧症有病者のうち服薬者※	185	62.3	0	0.0	1	16.7	4	23.5	18	48.6	65	63.7	97	72.9	123	59.1	80	67.2	61	75.3
正常高値血圧者	130	25.0	4	28.6	7	19.4	10	16.1	23	32.4	37	24.3	49	26.5	87	24.5	46	27.2	32	27.8
上記以外	93	17.9	8	57.1	23	63.9	35	56.5	11	15.5	13	8.6	3	1.6	60	16.9	4	2.4	2	1.7
総数	626	100.0	24	100.0	50	100.0	86	100.0	92	100.0	180	100.0	194	100.0	432	100.0	179	100.0	120	100.0
高血圧症有病者	272	43.5	0	0.0	1	2.0	6	7.0	32	34.8	100	55.6	133	68.6	182	42.1	108	60.3	89	74.2
(再掲)高血圧症有病者のうち服薬者※	190	69.9	0	0.0	0	0.0	2	33.3	16	50.0	60	60.0	112	84.2	114	62.6	73	67.6	76	85.4
正常高値血圧者	125	20.0	1	4.2	3	6.0	9	10.5	19	20.7	37	20.6	56	28.9	91	21.1	50	27.9	30	25.0
上記以外	229	36.6	23	95.8	46	92.0	71	82.6	41	44.6	43	23.9	5	2.6	159	36.8	21	11.7	1	0.8

注1)血圧を測定し、身体状況調査の問診において「血圧を下げる薬」の使用状況に回答した者を集計対象とした。

注2)判定の基準はp.11参照

注3)「服薬者」とは、質問票で「血圧を下げる薬」の使用有と回答した者

2-19 「脂質異常症が疑われる人」の状況(採血時間によらず、性・年齢階級別)

	総計		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総計	1,005	100.0	30	100.0	80	100.0	135	100.0	148	100.0	288	100.0	324	100.0	693	100.0	298	100.0	202	100.0
脂質異常症が疑われる人	313	31.1	2	6.7	9	11.3	18	13.3	38	25.7	98	34.0	148	45.7	215	31.0	123	41.3	87	43.1
(再掲)脂質異常症が疑われる人のうち服薬者※	196	62.6	1	50.0	1	11.1	5	27.8	21	55.3	66	67.3	102	68.9	130	60.5	83	67.5	64	73.6
上記以外	692	68.9	28	93.3	71	88.8	117	86.7	110	74.3	190	66.0	176	54.3	478	69.0	175	58.7	115	56.9
総計	462	100.0	11	100.0	35	100.0	57	100.0	64	100.0	135	100.0	160	100.0	317	100.0	150	100.0	99	100.0
脂質異常症が疑われる人	147	31.8	2	18.2	5	14.3	8	14.0	16	25.0	47	34.8	69	43.1	103	32.5	65	43.3	37	37.4
(再掲)脂質異常症が疑われる人のうち服薬者※	73	49.7	1	50.0	1	20.0	2	25.0	8	50.0	26	55.3	35	50.7	51	49.5	35	53.8	20	54.1
上記以外	315	68.2	9	81.8	30	85.7	49	86.0	48	75.0	88	65.2	91	56.9	214	67.5	85	56.7	62	62.6
総計	543	100.0	19	100.0	45	100.0	78	100.0	84	100.0	153	100.0	164	100.0	376	100.0	148	100.0	103	100.0
脂質異常症が疑われる人	166	30.6	0	0.0	4	8.9	10	12.8	22	26.2	51	33.3	79	48.2	112	29.8	58	39.2	50	48.5
(再掲)脂質異常症が疑われる人のうち服薬者※	123	74.1	0	0.0	0	0.0	3	30.0	13	59.1	40	78.4	67	84.8	79	70.5	48	82.8	44	88.0
上記以外	377	69.4	19	100.0	41	91.1	68	87.2	62	73.8	102	66.7	85	51.8	264	70.2	90	60.8	53	51.5

「脂質異常症が疑われる人」の判定
健康づくり国民調査では、空腹時採血が困難であるため、脂質異常症の診断基準項目である中性脂肪による判定は行わず、下記の通りとした。
HDLコレステロールが40mg/dl未満、もしくはコレステロールを下げる薬または中性脂肪(トリグリセライド)を下げる薬を服用している者。

2-20 運動習慣の状況(性・年齢階級別)

	総計		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 20-64歳		(再掲) 65-74歳以上		(再掲) 75歳以上	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	1,053	100.0	30	100.0	71	100.0	125	100.0	148	100.0	318	100.0	361	100.0	492	100.0	334	100.0	227	100.0
習慣無(健康上の理由)	42	4.0	1	3.3	0	0.0	2	1.6	4	2.7	14	4.4	21	5.8	10	2.0	15	4.5	17	7.5
習慣無(その他)	661	62.8	23	76.7	58	81.7	104	83.2	110	74.3	190	59.7	176	48.8	379	77.0	172	51.5	110	48.5
運動の習慣有	350	33.2	6	20.0	13	18.3	19	15.2	34	23.0	114	35.8	164	45.4	103	20.9	147	44.0	100	44.1
男性	475	100.0	9	100.0	30	100.0	53	100.0	64	100.0	145	100.0	174	100.0	205	100.0	159	100.0	111	100.0
習慣無(健康上の理由)	13	2.7	0	0.0	0	0.0	0	0.0	0	0.0	4	2.8	9	5.2	0	0.0	5	3.1	8	7.2
習慣無(その他)	289	60.8	7	77.8	22	73.3	45	84.9	46	71.9	86	59.3	83	47.7	157	76.6	86	54.1	46	41.4
運動の習慣有	173	36.4	2	22.2	8	26.7	8	15.1	18	28.1	55	37.9	82	47.1	48	23.4	68	42.8	57	51.4
女性	578	100.0	21	100.0	41	100.0	72	100.0	84	100.0	173	100.0	187	100.0	287	100.0	175	100.0	116	100.0
習慣無(健康上の理由)	29	5.0	1	4.8	0	0.0	2	2.8	4	4.8	10	5.8	12	6.4	10	3.5	10	5.7	9	7.8
習慣無(その他)	372	64.4	16	76.2	36	87.8	59	81.9	64	76.2	104	60.1	93	49.7	222	77.4	86	49.1	64	55.2
運動の習慣有	177	30.6	4	19.0	5	12.2	11	15.3	16	19.0	59	34.1	82	43.9	55	19.2	79	45.1	43	37.1

注)「運動の習慣有り」とは週2日以上30分以上の持続運動で1年以上継続している者

2-21 歩行数の平均値及び標準偏差(性・年齢階級別)

		人数	平均値(歩)	標準偏差
総数	総数	1,814	6,345	5,477
	15-19歳	67	7,208	3,717
	20-29歳	79	7,562	4,063
	30-39歳	194	6,925	5,618
	40-49歳	280	7,453	5,836
	50-59歳	294	7,040	6,067
	60-69歳	460	6,406	6,315
	70歳以上	440	4,504	3,473
	(再掲)20歳以上	1,747	6,312	5,532
	(再掲)20-64歳	1,053	7,085	5,905
	(再掲)65-74歳	441	5,901	5,234
	(再掲)75歳以上	253	3,809	3,086
男性	総数	820	6,764	5,708
	15-19歳	33	6,733	3,815
	20-29歳	33	9,203	4,624
	30-39歳	81	7,561	4,690
	40-49歳	130	8,284	7,597
	50-59歳	126	7,013	4,607
	60-69歳	207	6,783	6,787
	70歳以上	210	4,967	3,759
	(再掲)20歳以上	787	6,765	5,776
	(再掲)20-64歳	465	7,713	6,601
	(再掲)65-74歳	202	6,015	4,086
	(再掲)75歳以上	120	4,351	3,445
女性	総数	994	5,999	5,257
	15-19歳	34	7,670	3,616
	20-29歳	46	6,385	3,164
	30-39歳	113	6,470	6,178
	40-49歳	150	6,733	3,561
	50-59歳	168	7,060	6,978
	60-69歳	253	6,098	5,897
	70歳以上	230	4,082	3,139
	(再掲)20歳以上	960	5,940	5,297
	(再掲)20-64歳	588	6,588	5,242
	(再掲)65-74歳	239	5,804	6,044
	(再掲)75歳以上	133	3,319	2,641

2-22 歩行数の分布(性・年齢階級別)

	総数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 20歳以上		(再掲) 20-64歳		(再掲) 60歳以上		(再掲) 65-74歳		(再掲) 75歳以上		
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数
総数	1814	100.0	67	100.0	79	100.0	194	100.0	280	100.0	294	100.0	460	100.0	440	100.0	1,747	100.0	1,063	100.0	900	100.0	441	100.0	253	100.0	
2,000歩未満	218	12.0	3	4.5	3	3.8	12	6.2	12	4.3	24	8.2	49	10.7	115	26.1	215	12.3	73	6.9	164	18.2	55	12.5	87	34.4	
2,000-4,000歩未満	369	20.3	10	14.9	12	15.2	40	20.6	37	13.2	52	17.7	108	23.5	110	25.0	359	20.5	185	17.6	218	24.2	113	25.6	61	24.1	
4,000-6,000	422	23.3	14	20.9	12	15.2	47	24.2	79	28.2	68	23.1	105	22.8	97	22.0	408	23.4	253	24.0	202	22.4	99	22.4	56	22.1	
6,000-8,000	328	18.1	10	14.9	18	22.8	38	19.6	50	17.9	67	22.8	82	17.8	63	14.3	318	18.2	210	19.9	145	16.1	77	17.5	31	12.3	
8,000-10,000	204	11.2	14	20.9	21	26.6	22	11.3	41	14.6	33	11.2	47	10.2	26	5.9	190	10.9	140	13.3	73	8.1	43	9.8	7	2.8	
10,000歩以上	273	15.0	16	23.9	13	16.5	35	18.0	61	21.8	50	17.0	69	15.0	29	6.6	257	14.7	192	18.2	98	10.9	54	12.2	11	4.3	
0,000歩/10,000歩以上	392	21.6	24	35.8	26	32.9	49	25.3	82	29.3	72	24.5	95	20.7	44	10.0	368	21.1	275	26.1	139	15.4	79	17.9	14	5.5	
総数	820	100.0	33	100.0	33	100.0	81	100.0	130	100.0	126	100.0	207	100.0	210	100.0	787	100.0	465	100.0	417	100.0	202	100.0	120	100.0	
2,000歩未満	92	11.2	3	9.1	0	0.0	3	3.7	4	3.1	10	7.9	23	11.1	49	23.3	89	11.3	27	5.8	72	17.3	28	13.9	34	28.3	
2,000-4,000歩未満	143	17.4	6	18.2	2	6.1	14	17.3	13	10.0	20	15.9	44	21.3	44	21.0	137	17.4	73	15.7	88	21.1	37	18.3	27	22.5	
4,000-6,000	192	23.4	6	18.2	5	15.2	18	22.2	37	28.5	31	24.6	45	21.7	50	23.8	186	23.6	107	23.0	95	22.8	49	24.3	30	25.0	
6,000-8,000	146	17.8	2	6.1	7	21.2	17	21.0	22	16.9	29	23.0	34	16.4	35	16.7	144	18.3	87	18.7	69	16.5	39	19.3	18	15.0	
8,000-10,000	95	11.6	8	24.2	11	33.3	9	11.1	21	16.2	10	7.9	24	11.6	12	5.7	87	11.1	64	13.8	36	8.6	19	9.4	4	3.3	
10,000歩以上	152	18.5	8	24.2	8	24.2	20	24.7	33	25.4	26	20.6	37	17.9	20	9.5	144	18.3	107	23.0	57	13.7	30	14.9	7	5.8	
(再掲)2,000歩以上	181	22.1	12	36.4	12	36.4	24	29.6	38	29.2	30	23.8	44	21.3	21	10.0	169	21.5	128	27.5	65	15.6	34	16.8	7	5.8	
総数	994	100.0	34	100.0	46	100.0	113	100.0	150	100.0	168	100.0	253	100.0	230	100.0	960	100.0	588	100.0	483	100.0	239	100.0	133	100.0	
2,000歩未満	126	12.7	0	0.0	3	6.5	9	8.0	8	5.3	14	8.3	26	10.3	66	28.7	126	13.1	46	7.8	92	19.0	27	11.3	53	39.8	
2,000-4,000歩未満	226	22.7	4	11.8	10	21.7	26	23.0	24	16.0	32	19.0	64	25.3	66	28.7	222	23.1	112	19.0	130	26.9	76	31.8	34	25.6	
4,000-6,000	230	23.1	8	23.5	7	15.2	29	25.7	42	28.0	37	22.0	60	23.7	47	20.4	222	23.1	146	24.8	107	22.2	50	20.9	26	19.5	
6,000-8,000	182	18.3	8	23.5	11	23.9	21	18.6	28	18.7	38	22.6	48	19.0	28	12.2	174	18.1	123	20.9	76	15.7	38	15.9	13	9.8	
8,000-10,000	109	11.0	6	17.6	10	21.7	13	11.5	20	13.3	23	13.7	23	9.1	14	6.1	103	10.7	76	12.9	37	7.7	24	10.0	3	2.3	
10,000歩以上	121	12.2	8	23.5	5	10.9	15	13.3	28	18.7	24	14.3	32	12.6	9	3.9	113	11.8	85	14.5	41	8.5	24	10.0	4	3.0	
(再掲)8,300歩以上	211	21.2	12	35.3	14	30.4	25	22.1	44	29.3	42	25.0	51	20.2	23	10.0	199	20.7	147	25.0	74	15.3	45	18.8	7	5.3	